



ENT UK guideline on Necrotising Otitis Externa



References: 1. Mahdyoun P, Pulcini C, Gahide I, Raffaelli C, Savoldelli C, Castillo L, Guevara N. Necrotizing otitis externa: a systematic review. *Otol Neurotol.* 2013 Jun;34(4):620-9. 2. Courson AM, Vikram HR, Barrs DM. What are the criteria for terminating treatment for necrotizing (malignant) otitis externa? *Laryngoscope.* 2014 Feb;124(2):361-2. 3. Chawdhary G, Pankhania M, Douglas S, Bottrill I. Current management of necrotising otitis externa in the UK: survey of 221 UK otolaryngologists. *Acta Otolaryngol.* 2017 Aug;137(8):818-822. 4. Pankhania M, Bashyam A, Judd O, Jassar P. Antibiotic prescribing trends in necrotising otitis externa: a survey of 85 trusts in the United Kingdom: Our Experience. *Clin Otolaryngol.* 2016 Jun;41(3):293-6. 5. Phillips JS, Jones SE. Hyperbaric oxygen as an adjuvant treatment for malignant otitis externa. *Cochrane Database Syst Rev.* 2013 May 31;(5):CD004617. 6. Pulcini C, Mahdyoun P, Cua E, Gahide I, Castillo L, Guevara N. Antibiotic therapy in necrotising external otitis: case series of 32 patients and review of the literature. *Eur J Clin Microbiol Infect Dis.* 2012 Dec;31(12):3287-94. 7. Verim A, Naiboğlu B, Karaca Ç, Seneldir L, Kulekçi S, Oysu Ç. Clinical outcome parameters for necrotizing otitis externa. *Otol Neurotol.* 2014 Feb;35(2):371-6. Authors: Robert Nash (GOSH), Sarah Logan (UCLH), Trupti Patel (Imperial), Sherif Khalil (RNTNE), Shakeel Saeed (UCL)